

# Veterinary Dermatology

## Dermatologie vétérinaire

### Pemphigus foliaceus

Elizabeth Goodale

#### Introduction

**T**he epidermis consists of multiple layers of keratinocytes and makes up the outer layers of the skin (1). The keratinocytes in the epidermis are held together by desmosomes, which are cell-cell adhesion molecules made of several different proteins, such as desmocollin and desmoglein (1). These adhesion molecules are frequent targets for autoimmune diseases and their destruction results in superficial blistering or acantholysis (1). Pemphigus foliaceus is the most common autoimmune disease in dogs and is most often caused by autoantibodies targeting desmocollin-1 (2,3).

#### Clinical presentation

The classical lesions of pemphigus foliaceus are large pustules that span multiple hair follicles (1,4). The pustules often begin as papules and rapidly progress to crusts and erosions (1,4). Lesions occur most commonly on the trunk, inner pinnae, face, and footpads and are generally symmetrical (4,5). Mucosal lesions are extremely rare and are not considered a feature of pemphigus foliaceus (4). Pruritus is common and dogs with severe generalized disease can have anorexia, depression, fever, and weight loss (1,3,4).

Several breeds are reported as predisposed, but the Akita and chow chow are consistently reported as being overrepresented (3). Other breeds that have been reported as overrepresented include cocker spaniel, dachshund, doberman, collie, and Shar-pei (3). Secondary bacterial infections are common and can make the diagnosis more challenging (1,4). Involvement of the pinnae, nasal planum, and paw pads is more suggestive of pemphigus foliaceus than bacterial pyoderma which more commonly affects the ventrum (1).

Most cases of pemphigus foliaceus are idiopathic, but drug-induced cases have been reported, although the evidence is often weak (1,3,4). A history of atopic dermatitis and flea allergy dermatitis is common, but since these are often found in the general canine population it is unclear if there is truly an increased incidence (5,6). Disease progression can vary widely with some patients showing a rapid progression and other cases having a more gradual course (1,4).

#### Diagnostics

Cytology is a useful tool when initially examining a dog suspected to be suffering from pemphigus foliaceus. It is important to rule out bacterial pyoderma, which is the main

---

Dr. Goodale, Diplomate of the American College of Veterinary Dermatology, Guelph Veterinary Specialty Hospital, 1460 Gordon Street South, Guelph, Ontario N1L 1C8.

Address all correspondence to Dr. Elizabeth Goodale; e-mail: [egoodaledvm@gmail.com](mailto:egoodaledvm@gmail.com)

The Canadian Academy of Veterinary Dermatology (CAVD) is a not-for-profit organization that promotes veterinary dermatology in Canada and provides continuing education for veterinarians, animal health technicians/technologists and veterinary students. The CAVD welcomes applications for membership ([www.cavd.ca](http://www.cavd.ca)).

Use of this article is limited to a single copy for personal study. Anyone interested in obtaining reprints should contact the CVMA office ([hbroughton@cvma-acmv.org](mailto:hbroughton@cvma-acmv.org)) for additional copies or permission to use this material elsewhere.

differential diagnosis for pemphigus foliaceus (1). Cytology from cases of pemphigus foliaceus classically shows acantholytic keratinocytes associated with intact neutrophils and occasional eosinophils (1,4). Intracellular and extracellular bacteria may be seen if there is a secondary bacterial infection (4). Acantholytic keratinocytes are rounded, darkly staining nucleated keratinocytes with features of normally differentiated spinous or granular layer epithelial cells (1,3). Unfortunately, they are not seen exclusively with pemphigus foliaceus, as they have also been reported in cases of dermatophytosis and bacterial pyoderma so these must be ruled out (1).

A definitive diagnosis should be obtained using histopathology before instituting treatment. Dogs with secondary bacterial pyoderma should be treated with antibiotics for 2 to 4 wk before obtaining biopsies whenever possible. Biopsies should be taken from intact pustules or tightly adherent crusts. Care should be taken to avoid rupturing the pustules or removing the crusts while obtaining samples. This typically means that the skin should not be scrubbed. If a crust falls off the sample, it should be included in the formalin jar. Multiple punch or wedge biopsy samples should be taken and submitted to a dermatohistopathologist.

The classical histopathological findings for pemphigus foliaceus are subcorneal pustules containing rafts of acantholytic keratinocytes (7). These pustules are often layered and span multiple hair follicles (7). Nondegenerate neutrophils, and less commonly eosinophils, are typically present within the pustules (4,7). Special stains to rule out dermatophytosis should be considered (7).

Routine hematology and biochemistry should be performed before instituting therapy for pemphigus foliaceus (1). A severe neutrophilia is common (1).

## Treatment

Treatment of pemphigus foliaceus requires the use of immunosuppressive medications. Corticosteroids are the most commonly used medications either as a single agent or concurrently with a second immunosuppressive medication (1,4). Prednisone or prednisolone at a dose of 2 mg/kg body weight (BW) per day either given once daily or divided into twice daily dosing is the most common medication used (3). Some clinicians prefer to use methylprednisolone because it has less mineralocorticoid activity and may cause less polyuria and polydipsia (8). Refractory cases may respond to more potent glucocorticoids such as triamcinolone (0.2 to 0.6 mg/kg BW per day) or dexamethasone (0.2 to 0.4 mg/kg BW per day) (8). Short- and long-term side effects are common with corticosteroid usage and include polyuria, polydipsia, polyphagia, panting, increased risk of infections (urinary tract infections, pyoderma, demodicosis, dermatophytosis), alopecia, cutaneous atrophy, calcinosis cutis, and steroid hepatopathy (1,8,9).

One retrospective study did not find a significant difference in treatment outcomes for cases treated with corticosteroid monotherapy compared with prednisone and azathioprine used in combination (4). Regardless of these findings, many clinicians (including the author) typically begin with a combination of a corticosteroid and a second immunosuppressive medication

for cases of generalized pemphigus foliaceus and reserve single agent corticosteroids for mild cases (1).

If there is secondary bacterial infection it should be treated. Antibiotic use at the beginning of treatment was associated with a better outcome in 1 study, but other studies have not found a significant difference between cases treated with and without antibiotics (4,6).

Azathioprine is the most frequently reported secondary immunosuppressive medication used to treat pemphigus foliaceus (1,4). Azathioprine is a purine antagonist that is less expensive than many of the other immunosuppressive medications (1). Side effects include pancreatitis, bone marrow suppression, and hepatotoxicity, so a baseline complete blood cell count and serum biochemistry profile should be obtained prior to initiating therapy (1,10). This should be repeated after 2 wk, 4 wk, and monthly for the first 3 mo (10). Hepatotoxicity is most likely to occur within the first 2 to 4 wk and bone marrow suppression typically occurs with prolonged usage (10). It is typically dosed at 2.2 mg/kg BW once daily and clinical response can take 3 to 6 wk (1,9).

Cyclosporine has been reported to have variable success in managing pemphigus foliaceus (11–13). Cyclosporine is a calcineurin inhibitor that is commonly used to treat atopic dermatitis in dogs (1,13). Transient vomiting and diarrhea are the most common side effects and the maximal effect is typically seen after 4 wk (9,13). Anecdotally, freezing the capsules and administering them frozen reduces the incidence of vomiting and does not appear to impact the bioavailability (13,14). The use of the brand name microemulsified product is preferred, but this medication is unfortunately very expensive and can be cost-prohibitive in large dogs. Doses between 5 to 10 mg/kg BW per day are typically recommended for pemphigus foliaceus (8,13).

Mycophenolate mofetil is a purine antagonist that has not been used as extensively as azathioprine, but is gaining popularity in the treatment of pemphigus foliaceus (1,15). Mycophenolate mofetil has fewer side effects than azathioprine, so intensive monitoring of blood is not required (1,9). Diarrhea is the most common side effect reported and clinical response can take 3 to 8 wk (1,9,15). Doses of 20 to 40 mg/kg BW per day divided into 2 to 3 doses daily are recommended for pemphigus foliaceus (9,15).

Typically treatment with corticosteroids, with or without a secondary immunosuppressive agent, is instituted and continued until complete remission is achieved (1). The corticosteroids are then tapered by 25% every 2 to 4 wk until doses less than 0.5 mg/kg BW every other day for prednisone, 0.05 to 0.1 mg/kg BW every 2 to 3 d for dexamethasone, or 0.1 to 0.2 mg/kg BW every 2 to 3 d for triamcinolone are achieved (1). The secondary agent is then tapered with the ultimate goal of giving the corticosteroid and secondary agent every other day on alternating days (1,8).

For milder cases, the use of a combination of tetracycline (or doxycycline) and niacinamide has been reported (16). This combination of an antibiotic and a B vitamin has immunomodulatory effects, although the exact mechanisms are not fully understood (1). Hepatotoxicity has been rarely reported with doxycycline, and generally this combination is well-tolerated (9).

This drug combination has a very slow onset of action, which limits its utility in cases of pemphigus foliaceus that are generalized or severely pruritic (1). This can be combined with corticosteroids to try to hasten the onset of action (8). Tetracycline and niacinamide are dosed at 500 mg of each 3 times daily for dogs weighing > 10 kg and 250 mg of each 3 times daily for dogs weighing < 10 kg (8,16). Topical glucocorticoids or tacrolimus 0.1% could also be considered for localized lesions (1,8).

## Prognosis

Retrospective studies have found variable treatment success rates, but generally most cases have a positive response to treatment (1,3). An older retrospective study reported a 1-year survival of 53% and another had a case fatality rate of 60.5% (5,6). Newer retrospective studies have had more positive outcomes with 52% of cases achieving complete remission, 35% partial remission, and only 13% euthanized in 1 study (4). The most common reasons for euthanasia were treatment failure, treatment side effects, and poor quality of life (6). Survival beyond 10 mo is associated with a positive long-term outcome (6). There does not appear to be any difference in treatment outcomes for dogs with localized *versus* generalized disease or rapid onset *versus* slow onset (4). Most patients with pemphigus foliaceus will require lifelong treatment with medication although they are tapered to the lowest effective dose (1). A few patients may have prolonged periods of remission after their treatments have been withdrawn (4,17).

In conclusion, pemphigus foliaceus is the most common autoimmune disease affecting dogs (1). It is most commonly associated with bilaterally symmetrical crusting affecting the head, trunk, and footpads (4,5). A definitive diagnosis is made with histopathology, but cytology should be used to help differentiate cases from superficial pyoderma (1,4). Treatment typically requires immunosuppressive doses of glucocorticoids with or without a secondary immunosuppressive medication (1,4,5,8). Many clinicians initiate treatment with a combination of glucocorticoids and a secondary immunosuppressive medication but there is no clear evidence that this improves outcomes over glucocorticoid monotherapy (4). Most patients will require lifelong treatment, but the doses should be tapered to the lowest

effective dose (1). A few patients may be euthanized due to treatment failure, treatment side effects, or poor quality of life, which most commonly occurs during the early phase of treatment (6).

## References

1. Miller WH, Griffin CE, Campbell KL. Muller and Kirk's Small Animal Dermatology. 7th ed. St. Louis, Missouri: Elsevier, 2013.
2. Bizikova P, Dean GA, Hashimoto T, Olivry T. Cloning and establishment of canine desmocollin-1 as a major autoantigen in canine pemphigus foliaceus. *Vet Immunol Immunopathol* 2012;149:197–207.
3. Olivry T. A review of autoimmune skin diseases in domestic animals: I — Superficial pemphigus. *Vet Dermatol* 2006;17:291–305.
4. Mueller RS, Krebs I, Power HT, Fieseler KV. Pemphigus foliaceus in 91 dogs. *J Am Anim Hosp Assoc* 2006;42:189–196.
5. Ihrke PJ, Stannard A, Ardans A, Griffin CE. Pemphigus foliaceus in dogs: A review of 37 cases. *J Am Vet Med Assoc* 1985;186:59–66.
6. Gomez SM, Morris DO, Rosenbaum MR, Goldschmidt MH. Outcome and complications associated with treatment of pemphigus foliaceus in dogs: 43 cases (1994–2000). *J Am Vet Med Assoc* 2004;224:1312–1316.
7. Gross TL, Ihrke PJ, Walder EJ, Affolter VK. Skin diseases of the dog and cat: Clinical and histopathologic diagnosis. 2nd ed. Ames, Iowa: Blackwell Science; 2005.
8. Rosenkrantz WS. Pemphigus: Current therapy. *Vet Dermatol* 2004;15:90–98.
9. Plumb DC. Plumb's Veterinary Drug Handbook. 7th ed. Stockholm, Wisconsin: PharmaVet; 2011.
10. Wallisch K, Trepanier LA. Incidence, timing, and risk factors of azathioprine hepatotoxicosis in dogs. *J Vet Intern Med* 2015;29:513–518.
11. Olivry T, Rivierre C, Murphy KM. Efficacy of cyclosporine for treatment induction of canine pemphigus foliaceus. *Vet Rec* 2003;152:53–54.
12. Kim JH, Sur JH, Park C, Yoo JH, et al. Therapeutic administration of cyclosporine in a dog with pemphigus foliaceus. *J Vet Clin* 2009;26:340–343.
13. Forsythe P, Paterson S. Cyclosporin 10 years on: Indications and efficacy. *Vet Rec* 2014;174:13–21.
14. Bachtel JC, Pendergraft JS, Rosychuk RA, Gustafson DL, Hansen RJ, Lunghofer PJ. Comparison of the stability and pharmacokinetics in dogs of modified cyclosporin capsules stored at –20°C and room temperature. *Vet Dermatol* 2015;26:228–e50.
15. Ackermann AL, May ER, Frank LA. Use of mycophenolate mofetil to treat immune-mediated skin disease in 14 dogs — A retrospective evaluation. *Vet Dermatol* 2017;28:195–e44.
16. White SD, Rosychuk RA, Reinke SI, Paradis M. Use of tetracycline and niacinamide for treatment of autoimmune skin disease in 31 dogs. *J Am Vet Med Assoc* 1992;200:1497–1500.
17. Olivry T, Bergvall KE, Atlee B. Prolonged remission after immunosuppressive therapy in six dogs with pemphigus foliaceus. *Vet Dermatol* 2004;15:245–252.